Math. Struct. in Comp. Science (2000), *vol.* 10, *pp.* 137–163. Printed in the United Kingdom
© 2000 Cambridge University Press

Unique factorisation lifting functors and categories of linearly-controlled processes

MARTA BUNGE[†] and MARCELO P. FIORE[‡]

[†] *Department of Mathematics and Statistics, McGill University, 805 Sherbrooke Street West, Montreal, Quebec, Canada H3A 2K6.*

[‡] *School of Cognitive and Computing Sciences, University of Sussex, Falmer, Brighton, BN1 9QH, UK.*

Received 16 November 1998; revised 10 September 1999

Dedicated to Jim Lambek on the occasion of his 75th birthday

We consider processes consisting of a category of states varying over a control category as prescribed by a unique factorisation lifting functor. After a brief analysis of the structure of general processes in this setting, we restrict attention to linearly-controlled ones. To this end, we introduce and study a notion of path-linearisable category in which any two paths of morphisms with equal composites can be linearised (or interleaved) in a canonical fashion. Our main result is that categories of linearly-controlled processes (*viz*., processes controlled by path-linearisable categories) are sheaf models.

1. Introduction

This work is an investigation into the mathematical structure of processes. The processes to be considered embody a notion of state space varying according to a control. This we formalise as a category of states (and their inter-relations) $\mathbb{X}$ equipped with a control functor $f \downarrow \begin{matrix}\mathbb{X}\\ \\ \mathbb{C}\end{matrix}$.

There are different ways in which the control category $\mathbb{C}$ may be required to control the state category $\mathbb{X}$; these depend, of course, on the properties required from the control functor f. For instance, one may require the control functor to be a discrete opfibration or, dually, a discrete fibration. The first requirement corresponds to imposing a determinacy property on the state category (Lawvere 1986b; Lawvere 1986c), as for every state X in $\mathbb{X}$ and every control path $\gamma : f(X) \longrightarrow C$ there should exist a unique evolution $e : X \longrightarrow Y$ in $\mathbb{X}$ along γ (*i.e*., such that $f(e) = \gamma$). The second case corresponds essentially to the

[†] Research supported by grants from the Natural Sciences and Engineering Council of Canada and from the Ministére de l'Education du Québec.

[‡] Research done while visiting the Centre des Recherches en Théorie des Catégories at McGill University during September–October 1998.

presheaf models of Winskel *et al.* (Joyal *et al.* 1996; Cattani and Winskel 1997; Winskel and Nielsen 1997). It allows non-determinism but, as we explain in Section 2, imposes a particular notion of state on the state category. The approach adopted here will be to consider a more general notion of control functor; *viz.*, *unique factorisation lifting (ufl) functors*. Our motivation for doing so comes from computer science where ufl functors can be used to provide a conceptual understanding of certain models for concurrency that have been dealt with in an *ad hoc* manner (see Sections 2 and 4). For motivations relevant to continuum physics see Lawvere (1986b) and Lawvere (1986c).

Intuitively, in a ufl functor $f\downarrow^{\mathbb{X}}_{\mathbb{C}}$, every evolution $e : X \longrightarrow Y$ in $\mathbb{X}$ is controlled by $\mathbb{C}$ in that whenever the control path $f(e)$ in $\mathbb{C}$ may be performed in two stages (γ_0, γ_1) (that is, $f(e) = \gamma_0 \cdot \gamma_1$), the evolution e uniquely decomposes as two evolutions (e_0, e_1) (that is, $e = e_0 \cdot e_1$) taking place along γ_0 and γ_1, respectively (that is, $f(e_0) = \gamma_0$ and $f(e_1) = \gamma_1$).

Thus, a first rough notion of category of processes can be taken to be the category $\mathbf{UFL}/\mathbb{C}$, where **UFL** is the subcategory of **Cat** of ufl functors. We show in Section 3 that $\mathbf{UFL}/\mathbb{C}$ may be embedded (as a full reflective subcategory) in a presheaf topos; *viz.*, $\widehat{\mathbb{C}_{\#}}$ where $\mathbb{C}_{\#}$ is Mac Lane's twisted arrow category (Mac Lane 1971). On the one hand, this result establishes connections with presheaf models; thus opening the possibility of reusing the techniques developed in that setting (Cattani and Winskel 1997). On the other hand, it highlights important differences. First, the path category $\mathbb{C}_{\#}$ is obtained from the control category $\mathbb{C}$ and this passage equips the former with extra structure; notably notions of cover. Second, presheaf models can be over generous. For instance, by considering *all* presheaves in $\widehat{\mathbb{C}_{\#}}$ as processes, one is doomed to rule out important notions of state (*e.g.*, as in transition systems; see Section 2). This may be overcome by restricting to presheaves in $\widehat{\mathbb{C}_{\#}}$ satisfying certain orthogonality (sheaf) conditions; for these are in bijective correspondence with ufl functors over $\mathbb{C}$ (the bijection extracts a state category out of a presheaf). However, this mathematical equivalence should not undermine the fibrational viewpoint provided by ufl functors.

Finally, to establish the categories of processes that will be the subject study of this paper we need to analyse the nature of control categories. Various interpretations for them are possible. For example, one may regard control categories as providing time duration for the evolutions of a process; this is the viewpoint adopted in Lawvere (1986b) and Lawvere (1986c). Natural examples under this interpretation are the additive monoid of natural numbers (discrete time) and the additive monoid of non-negative real numbers (continuous time). Another view of control categories is as providing observations (*e.g.*, time passing or actions being performed); this is the interpretation explored here. Our basic example is the free monoid on a set of actions. In particular, when the set of actions is a singleton (containing a 'tic') we obtain the monoid of discrete time. The relevant structure of these and similar examples of control categories is axiomatised as so-called *path-linearisable* categories, which we introduce and study in Section 5. Informally, path-linearisable control categories are those in which any two control paths of morphisms with equal composites can be linearised (or interleaved) in a canonical fashion; see diagram (18). This property is better defined as the conjunction of a 'cancellation'

and a 'fill-in' property (see Definition 5.1). The cancellation property has been considered by Lawvere in the context of processes in Lawvere (1989) and, as argued there, seems to be inherent in the notion of a given evolution of a process. The further imposition of the fill-in property has important conceptual and mathematical consequences. Conceptually, the consideration of processes over a path-linearisable control category corresponds to restricting attention to processes that are linearly controlled; that is, to interleaving models, in the jargon of computer science. From the mathematical perspective, as shown in Section 5, the cancellation and fill-in properties on a category $\mathbb{C}$ are equivalent to an 'interval glueing' property used in Bunge and Niefield (1998) to show that $\mathbf{UFL}/\mathbb{C}$ is a (model-generated) topos. We give another proof of this result in Section 6 where we show that, for $\mathbb{C}$ a path-linearisable category, $\mathbf{UFL}/\mathbb{C}$ is a topos (of sheaves on $\mathbb{C}_{\#}$). We remark that Johnstone showed that for $\mathbb{C}$ the commutative square, the simplest example of failure of fill-in, the category $\mathbf{UFL}/\mathbb{C}$ is not a topos. This result can be found in Johnstone (1991), where it is also shown that, for a category $\mathbb{C}$ satisfying a *factorisation preordered* (our cancellation) and a *factorisation strongly connected* (our fill-in) condition, $\mathbf{UFL}/\mathbb{C}$ is a sheaf topos.

Organisation of the paper. We start in Section 2 with background material motivated from the point of view of computer science. In Section 3, the category **UFL** is defined and its basic properties are studied. Section 4 explores applications of ufl functors to concurrency. In particular, we show that they may be used in the study of interleaving models like transition systems. In Section 5, we introduce *path-linearisable* categories. Our main use for them is in Section 6 where, for $\mathbb{C}$ a path-linearisable category, we exhibit the category $\mathbf{UFL}/\mathbb{C}$ as a sheaf topos. These toposes may be regarded as models of linearly-controlled processes. Some concluding remarks are provided in Section 7.

2. Background and motivation

Labelled transition systems are a model of computation widely used in concurrency (Winskel and Nielsen 1995). We recall the basic definitions.

A *transition system* over a set of *actions* A consists of a set of *states* S and a transition relation $\longrightarrow \subseteq S \times A \times S$. A tuple $(s, a, s') \in \longrightarrow$ is usually denoted as $s \xrightarrow{a} s'$; its intended meaning is that at the state s the system may perform the action a and, in doing so, evolve to the state s'. The category of transition systems over a set of actions A, denoted $\mathbf{TS}_A$, has transition systems over A as objects and morphisms $\sigma : (S, \rightarrow) \longrightarrow (S', \rightarrow')$ where $\sigma : S \longrightarrow S'$ is a function such that $s \xrightarrow{a} s'$ implies $\sigma(s) \xrightarrow{a} \sigma(s')$.

One is sometimes interested in transition systems with an *initial state* and morphisms preserving it. This category is obtained as the slice $\mathbf{TS}_A^{\bullet} = U/\mathbf{TS}_A$ where $U = \boxed{\bullet}$ is the transition system with one state and no transitions.

An important subclass of transition systems is that of *synchronisation trees* (Winskel and Nielsen 1995). Roughly, these are transition systems with a distinguished initial state, with no cycles, and such that

$$s_1 \xrightarrow{a_1} s \text{ and } s_2 \xrightarrow{a_2} s \text{ imply } s_1 = s_2 \text{ and } a_1 = a_2 \ . \tag{1}$$

A conceptual description of the category of synchronisation trees is as certain *presheaf models* (Joyal *et al.* 1996; Cattani and Winskel 1997; Winskel and Nielsen 1997). We present this viewpoint next.

For a set A, let $\mathbf{P}(A)$ be the poset $(A^*, \leqslant)$ of finite sequences over A ordered by prefix. The category of synchronisation trees over a set of actions A is $\mathbf{ST}_A = \widetilde{\mathbf{P}(A)}$, where for a small category $\mathbb{P}$ with an initial object we write $\widetilde{\mathbb{P}}$ for the full subcategory of $\widehat{\mathbb{P}} = [\mathbb{P}^{\mathrm{op}}, \mathbf{Set}]$ of *rooted* presheaves. These are presheaves $T : \mathbb{P}^{\mathrm{op}} \longrightarrow \mathbf{Set}$ satisfying either of the following equivalent conditions: for $\perp$ an initial object in $\mathbb{P}$,

(R–*i*) The set $T(\perp)$ is terminal in **Set**.

(R–*ii*) There exists a unique map $\perp \longrightarrow T$ in $\widehat{\mathbb{P}}$ (that is, $0 \longrightarrow \perp$ is *orthogonal* (Freyd and Kelly 1972) to $T \longrightarrow 1$ in $\widehat{\mathbb{P}}$).

In the definition of the category of synchronisation trees, we have implicitly identified the rooted presheaf $T : \mathbf{P}(A)^{\mathrm{op}} \longrightarrow \mathbf{Set}$ with the synchronisation tree with initial state given by the unique element of $T(\varepsilon)$; set of states given by $\sum_{\alpha \in A^*} T(\alpha)$; and transition relation given by

$$(\alpha, s) \xrightarrow{a} (\alpha', s') \quad \text{iff} \quad \alpha' = \alpha \cdot a \text{ and } s = T(\alpha \leqslant \alpha')(s') \ .$$

This construction is the well-known *Grothendieck construction* (Mac Lane and Moerdijk 1994; Borceux 1994) in disguise. The Grothendieck construction applies to presheaf categories over an arbitrary small category $\mathbb{P}$, and provides a *fibrational* view of $\widehat{\mathbb{P}}$ as a full subcategory of $\mathbf{Cat}/\mathbb{P}$. It associates a presheaf $T : \mathbb{P}^{\mathrm{op}} \longrightarrow \mathbf{Set}$ with the functor $\begin{smallmatrix} \int T \\ \downarrow \\ \mathbb{P} \end{smallmatrix}$ given by an obvious projection, where the *category of elements* $\int T$ has set of objects given by $\sum_{\alpha \in \mathbb{P}} T(\alpha)$; morphisms $(\alpha, s) \xrightarrow{a} (\alpha', s')$ given by morphisms $a : \alpha \longrightarrow \alpha'$ in $\mathbb{P}$ such that $s = T(a)(s')$; identities given by $\mathrm{id}_{(\alpha,s)} = \mathrm{id}_\alpha$; and composition given as in $\mathbb{P}$. This mapping extends to an embedding

$$\widehat{\mathbb{P}} \hookrightarrow \mathbf{Cat}/\mathbb{P} \tag{2}$$

whose image consists of so-called *discrete fibrations*. Moreover, in the presence of the axiom of choice, the embedding (2) cuts down to an equivalence

$$\widehat{\mathbb{P}} \ \simeq \ \mathbf{DFib}/\mathbb{P} \tag{3}$$

where **DFib** denotes the subcategory of **Cat** of discrete fibrations. Recall that a functor $\begin{smallmatrix} \mathbb{T} \\ t\downarrow \\ \mathbb{P} \end{smallmatrix}$ is said to be a *discrete fibration* whenever it satisfies the following property.

(DF) In the situation

$$\begin{array}{ccc} & & s' \\ & & \vdots\, t \\ \alpha & \xrightarrow[a]{} & \alpha' \end{array} \tag{4}$$

there exists a unique morphism $s \longrightarrow s'$ in $\mathbb{T}$ such that

$$\begin{array}{ccc} s & \longrightarrow & s' \\ & \exists! \; \vdots & t \\ \alpha & \longrightarrow & \alpha' \end{array} \tag{5}$$

Thus, the property (DF) establishes that every commutative square

$$\begin{array}{ccc} \mathbf{1} & \longrightarrow & \mathbb{T} \\ {\scriptstyle (1)}\downarrow & \nearrow_{\exists!} & \downarrow{\scriptstyle t} \\ \mathbf{2} & \longrightarrow & \mathbb{P} \end{array}$$

has a unique fill-in as indicated (*i.e.*, that $(1) : \mathbf{1} \longrightarrow \mathbf{2}$ is orthogonal to $t : \mathbb{T} \longrightarrow \mathbb{P}$ in **Cat**). Dually, a functor is a *discrete opfibration* if $(0) : \mathbf{1} \longrightarrow \mathbf{2}$ is orthogonal to it.

The equivalence (3) associates to a discrete fibration $\begin{array}{c}\mathbb{T}\\ t\downarrow\\ \mathbb{P}\end{array}$ the presheaf $T : \mathbb{P}^{\mathrm{op}} \longrightarrow \mathbf{Set}$ where $T(\alpha) = \{ s \in \mathbb{T}_0 \mid t(s) = \alpha \}$ and, for $a : \alpha \longrightarrow \alpha'$ in $\mathbb{P}$, the function $T(a) : T(\alpha') \longrightarrow T(\alpha)$ maps an s' as in (4) to the unique s as in (5).

Observe that from the fibrational perspective the rootness conditions (R) above can be expressed as a *unique lifting* property. Indeed, for a small category $\mathbb{P}$ with an initial object $\perp$, we have the equivalence

$$\widetilde{\mathbb{P}} \;\simeq\; \mathbf{DFib}_0/\mathbb{P}$$

where $\mathbf{DFib}_0$ is the full subcategory of **DFib** consisting of discrete fibrations $\begin{array}{c}\mathbb{T}\\ t\downarrow\\ \mathbb{P}\end{array}$ such that $(\perp) : \mathbf{1} \longrightarrow \mathbb{P}$ has a unique lifting along t as follows

$$\begin{array}{ccc} & & \mathbb{T} \\ & \nearrow_{\exists!} & \downarrow{\scriptstyle t} \\ \mathbf{1} & \xrightarrow[(\perp)]{} & \mathbb{P} \end{array} \tag{6}$$

Since t is a discrete fibration, the object provided by the lifting $\mathbf{1} \longrightarrow \mathbb{T}$ of (6) is initial in $\mathbb{T}$; hence t creates initial objects.

Following the work on presheaf models, the category $\mathbf{DFib}/\mathbb{P}$ may be regarded as a category of *processes* over the category of *computation paths* $\mathbb{P}$. Indeed, a (discrete) fibration $\begin{array}{c}\mathbb{T}\\ t\downarrow\\ \mathbb{P}\end{array}$ embodies a category of states and transitions (the objects and morphisms of $\mathbb{T}$) varying along *observations* (the morphisms of $\mathbb{P}$) as specified by the *labelling* functor t. More concretely, a morphism $s \xrightarrow{e} s'$ in $\mathbb{T}$ witnesses the capability of the system to evolve from the state s to the state s' producing the observable behaviour $t(e)$ (Winskel

and Nielsen 1997). In this light, the property (DF) has the following computational interpretation: a state s' occurring after an observation a, as indicated in (4) above, comes from a unique state s and transition $s \longrightarrow s'$ along which a is observed. Notice that for discrete fibrations (equivalently presheaves) the notion of state captured by this interpretation is that of the *computation state* associated to the behaviour of a system, as opposed to the states in which a system can be. Indeed, we have the following property:

$$\text{for } s_1 \xrightarrow{e_1} s \text{ and } s_2 \xrightarrow{e_2} s, \text{ if } t(e_1) = t(e_2) \text{ then } s_1 = s_2 \text{ and } e_1 = e_2$$

which is analogous to property (1) of synchronisation trees.

We will show below that, by broadening the notion of fibration, one can develop a theory for transition systems with a treatment of bisimulation via open-maps (Joyal *et al.* 1996) in the same spirit as the presheaf models of Winskel's *et al.*. The required fibration condition amounts to a *unique factorisation lifting* property (Lawvere 1986b; Street 1996; Bunge and Niefield 1998). We will illustrate this concept in the context of transition systems.

Let $(S, \longrightarrow)$ be an A-labelled transition system and let $\mathscr{F}(S, \longrightarrow)$ be the free category on its underlying graph (that is, its reflexive-transitive closure). Explicitly, the small category $\mathscr{F}(S, \longrightarrow)$ has set of objects given by S; morphisms $s \longrightarrow s'$ given by finite *paths* (or *runs*, or *histories*) $[s \xrightarrow{a_1} s_1 \longrightarrow \cdots \xrightarrow{a_n} s']$ of the system; identities given by the empty paths $[s]$; and composition given by concatenation of paths. Writing $\mathbf{M}(A)$ for the free monoid on A, we have a projection functor $f \downarrow^{\mathscr{F}(S,\longrightarrow)}_{\mathbf{M}(A)}$, mapping a path $[s \xrightarrow{a_1} s_1 \longrightarrow \cdots \xrightarrow{a_n} s']$ to the sequence of actions $a_1 \cdots a_n$ performed along it, satisfying the following fibrational *unique factorisation lifting (ufl)* property: for every $e = [s \xrightarrow{a_1} s_1 \longrightarrow \cdots \xrightarrow{a_n} s']$ in $\mathscr{F}(S, \longrightarrow)$ and every factorisation $(\alpha_0, \alpha_1) = (a_1 \cdots a_i, a_{i+1} \cdots a_n)$ of $f(e) = a_1 \cdots a_i a_{i+1} \cdots a_n$ in $\mathbf{M}(A)$ there is a unique factorisation (e_0, e_1) of e in $\mathscr{F}(S, \longrightarrow)$ (namely $([s \xrightarrow{a_1} s_1 \longrightarrow \cdots \xrightarrow{a_i} s_i], [s_i \xrightarrow{a_{i+1}} s_{i+1} \longrightarrow \cdots \xrightarrow{a_n} s'])$) such that $f(\alpha_0) = e_0$ and $f(\alpha_1) = e_1$.

In Section 4, the category

$$\mathbf{UFL}/\mathbf{M}(A),$$


where **UFL** (Bunge and Niefield 1998) denotes the subcategory of **Cat** consisting of ufl functors (see Definition 3.1), will be studied as the category of models of computation given by *transition graphs*.

3. Unique factorisation lifting functors


We introduce unique factorisation lifting (ufl) functors (Lawvere 1986b; Street 1996; Bunge and Niefield 1998) and study their basic properties.

Definition 3.1. A *ufl functor* is a functor $\mathbb{X} \longrightarrow \mathbb{B}$ satisfying the following fibrational unique factorisation lifting property.


(UFL) In the situation



there exists unique $A \longrightarrow B$ and $B \longrightarrow C$ such that



Thus, the property (UFL) establishes that every commutative square



has a unique fill-in as indicated (*i.e.*, that $(0,2) : \mathbf{2} \longrightarrow \mathbf{3}$ is orthogonal to $\mathbb{X} \longrightarrow \mathbb{B}$ in **Cat**).

Since ufl functors are closed under identities and composition, they form a subcategory of **Cat**, which we denote **UFL**. Moreover, since whenever $\mathbb{X} \longrightarrow \mathbb{Y} \longrightarrow \mathbb{B}$ and $\mathbb{Y} \longrightarrow \mathbb{B}$ are ufl functors, so is $\mathbb{X} \longrightarrow \mathbb{Y}$, we have that $\mathbf{UFL}/\mathbb{B}$ is a full subcategory of $\mathbf{Cat}/\mathbb{B}$.

As can be easily seen, ufl functors reflect identities.

Lemma 3.2. For a ufl functor f, if $f(x) = \mathrm{id}$, then $x = \mathrm{id}$.

We consider examples of ufl functors.


Example 3.3.

1 Discrete fibrations and discrete opfibrations are ufl functors.

2 The *difference* functor $\Delta_A : \mathbf{P}(A) \longrightarrow \mathbf{M}(A)$ mapping $(\alpha \leqslant \alpha \cdot \alpha')$ to α' is a ufl functor. In particular, for $A = 1$, we obtain the usual difference functor $(m \leqslant n) \longmapsto n - m$ from $\boldsymbol{\omega} = \mathbf{P}(1)$, the poset of natural numbers, to $\mathbf{N} = \mathbf{M}(1)$, the additive monoid of natural numbers.

3 We let **Graph** (Lawvere 1986a; Lawvere 1989) be the presheaf category of *graphs* $\widehat{\mathbf{G}}$ where $\mathbf{G}^{\mathrm{op}} = \boxed{\mathsf{N} \;\overleftarrow{\overleftarrow{}}\; \mathsf{E}}$, and we write $\mathscr{F}$ for the left adjoint to the forgetful functor $\mathbf{Cat} \longrightarrow \mathbf{Graph} : \mathbb{C} \longmapsto (\mathbb{C}_0 \;\overset{\overset{\mathrm{dom}}{\longleftarrow}}{\underset{\mathrm{cod}}{\longleftarrow}}\; \mathbb{C}_1)$ providing the free category on a graph.

Consider the following *loop* construction ℓ defined by the adjoint situation below



Explicitly, for a set A, the graph ℓA is given by $(1 \underset{\longleftarrow}{\overset{\longleftarrow}{\longleftarrow}} A)$.
We have that $\ell 1$ is the terminal graph and that $\mathscr{F}(\ell 1) \cong \mathbf{N}$. Moreover, for a graph G, the unique graph homomorphism $G \longrightarrow \ell 1$ induces a *length* functor $\mathscr{F}G \longrightarrow \mathbf{N}$ that is a ufl functor (Schanuel and Street 1988; Street 1996).
More generally, for a set A, we have that $\mathscr{F}(\ell A) \cong \mathbf{M}(A)$ and that a graph G together with a *labelling* homomorphism $G \longrightarrow \ell A$ induces a ufl functor $\mathscr{F}G \longrightarrow \mathbf{M}(A)$.

The rest of the section is devoted to the study of the slice categories $\mathbf{UFL}/\mathbb{B}$. These we characterise as certain full subcategories of a presheaf category by means of orthogonality conditions (Lemma 3.7). Since the representables are ufl functors, we obtain a Yoneda embedding for $\mathbf{UFL}/\mathbb{B}$.

The required category of representables is Mac Lane's 'twisted arrow category' (see Mac Lane (1971, page 223)) of the base category. We recall the construction.

Definition 3.4. For a small category $\mathbb{B}$ let $\mathbb{B}_{\#}$ be the small category with objects given by the morphisms of $\mathbb{B}$, arrows $\alpha \longrightarrow \beta$ given by pairs of morphisms (φ, γ) in $\mathbb{B}$ such that

$$\begin{array}{ccc} A & \xleftarrow{\varphi} & B \\ {\scriptstyle\alpha}\downarrow & & \downarrow{\scriptstyle\beta} \\ A' & \xrightarrow[\gamma]{} & B' \end{array}$$

commutes, identities given by pairs of identities in $\mathbb{B}$, and composition given pointwise.

Example 3.5. The twisted arrow category $\mathbf{M}(A)_{\#}$ has objects given by finite sequences $\alpha \in A^*$ and morphisms given by *extensions* $(\varphi, \gamma) : \alpha \longrightarrow \varphi \cdot \alpha \cdot \gamma$ $(\alpha, \varphi, \gamma \in A^*)$. Notice that the poset $\mathbf{P}(A)$ appears as the subcategory of $\mathbf{M}(A)_{\#}$ consisting of the *post-extensions* $(\varepsilon, \gamma) : \alpha \longrightarrow \alpha \cdot \gamma$ $(\alpha, \gamma \in A^*)$, whilst the dual poset $\mathbf{P}(A)^{\mathrm{op}}$ appears as the subcategory of $\mathbf{M}(A)_{\#}$ of the *pre-extensions* $(\varphi, \varepsilon) : \alpha \longrightarrow \varphi \cdot \alpha$ $(\alpha, \varphi \in A^*)$. Thus, we may regard $\mathbf{M}(A)_{\#}$ as a *path* category extending $\mathbf{P}(A)$ with intertwined past and future (or back and forth) observations.

The mapping $(_)_{\#} : \mathbb{B} \longmapsto \mathbb{B}_{\#}$ extends to an endofunctor on **Cat** in an obvious way. It was show in Bunge and Niefield (1998) that,

$$f \text{ is a ufl functor iff } f_{\#} \text{ is a discrete fibration.}$$

Thus we obtain functors $\mathbf{UFL}/\mathbb{B} \longrightarrow \mathbf{DFib}/\mathbb{B}_{\#}$, which we also denote $(_)_{\#}$.

Proposition 3.6. For $g \underset{\mathbb{B}}{\overset{\mathbb{Y}}{\downarrow}}, f \underset{\mathbb{B}}{\overset{\mathbb{X}}{\downarrow}} \in |\mathbf{Cat}/\mathbb{B}|$, consider $H : g_{\#} \longrightarrow f_{\#}$ in $\mathbf{Cat}/\mathbb{B}_{\#}$. If f is a ufl functor, there exists a (necessarily unique) functor $h : \mathbb{Y} \longrightarrow \mathbb{X}$ such that $h : g \longrightarrow f$ in $\mathbf{Cat}/\mathbb{B}$ and $H = h_{\#}$.

Proof. The functor H has the following properties:

— For $Y \in \mathbb{Y}_0$, we have that $H(\mathrm{id}_Y) = \mathrm{id}_X$ for some (necessarily unique) $X \in \mathbb{X}_0$, which we set to be $h(Y)$.

This follows from Lemma 3.2, because $f(H(\mathrm{id}_Y)) = \mathrm{id}_{gY}$ using the fact that $f_\# \circ H = g_\#$ and the definition of $(_)_\#$.

— For $y : Y \longrightarrow Y'$ in $\mathbb{Y}$,

1 $\mathrm{dom}(Hy) = h(Y)$ and $H(\mathrm{id}_Y, y) = (\mathrm{id}_{hY}, Hy)$.

2 $\mathrm{cod}(Hy) = h(Y')$ and $H(y, \mathrm{id}_{Y'}) = (Hy, \mathrm{id}_{hY'})$.

w We only show (1), as (2) is proved similarly. Consider $(\mathrm{id}_Y, y) : \mathrm{id}_Y \longrightarrow y$ in $\mathbb{Y}_\#$, and let

$$H(\mathrm{id}_Y, y) \quad = \quad \begin{array}{ccc} h(Y) & \xleftarrow{x_0} & X_0 \\ {\scriptstyle H(\mathrm{id}_Y)=\mathrm{id}_{hY}}\downarrow & & \downarrow{\scriptstyle H(y)} \\ h(Y) & \xrightarrow[x_1]{} & X_1 \end{array}$$

Using the fact that $f_\# \circ H = g_\#$ and the definition of $(_)_\#$, we have $(f(x_0), f(x_1)) = (\mathrm{id}_{gY}, g(y))$. Thus, by Lemma 3.2, we have that $X_0 = h(Y)$ and $x_0 = \mathrm{id}_{hY}$, and so also that $x_1 = H(y)$.

From the above we obtain the following two further properties:

— For $Y \xrightarrow{y} Y' \xrightarrow{y'} Y''$ in $\mathbb{Y}$,

$$H(y \cdot y') \quad = \quad H(y) \cdot H(y').$$

— For $(y_0, y_1) : y \longrightarrow y'$ in $\mathbb{Y}_\#$,

$$H(y_0, y_1) \quad = \quad (Hy_0, Hy_1).$$

Finally, defining $h : \mathbb{Y} \longrightarrow \mathbb{X} : y \longmapsto H(y)$, we are done. □

It follows from this proposition that the functor $(_)_\# : \mathbf{UFL}/\mathbb{B} \longrightarrow \mathbf{DFib}/\mathbb{B}_\#$ is an embedding. We proceed to characterise its image.

Lemma 3.7. For $p \in |\mathbf{DFib}/\mathbb{B}_\#|$ the following are equivalent.

1 For some $f \in |\mathbf{UFL}/\mathbb{B}|$, the discrete fibration p and $f_\#$ are isomorphic.


2 The discrete fibration p creates the following pushouts

$$\begin{array}{ccc} & \beta & \\ {\scriptstyle (\alpha,\mathrm{id}_C)}\swarrow & & \searrow{\scriptstyle (\mathrm{id}_B,\gamma)} \\ \alpha \cdot \beta & & \beta \cdot \gamma \\ {\scriptstyle (\mathrm{id}_A,\gamma)}\searrow & & \swarrow{\scriptstyle (\alpha,\mathrm{id}_D)} \\ & \alpha \cdot \beta \cdot \gamma & \end{array} \qquad (A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D \text{ in } \mathbb{B}) \tag{7}$$

(that is, the pushouts (7) have a unique lifting along p).

3 For P the presheaf in $\widehat{\mathbb{B}_\#}$ associated to the discrete fibration p in $\mathbf{DFib}/\mathbb{B}_\#$, the


diagram



$(A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D \text{ in } \mathbb{B})$ (8)


is a pullback in **Set**.

Proof.
(1) $\Longrightarrow$ (2) For $f \in |\mathbf{UFL}/\mathbb{B}|$ we show that $f_\#$ creates the pushouts (7). Indeed, assume




(9)


Then, since f is a ufl functor, it follows from Lemma 3.2 that $u_1 = \mathrm{id}$ and $w_0 = \mathrm{id}$; and hence that $u = u_0 \cdot v$ and $w = v \cdot w_1$. Thus, (9) has as unique lifting the pushout



(2) $\Longrightarrow$ (3) Because (8) is a pullback iff in the situation



there exists a unique $\overset{x}{\underset{\alpha\cdot\beta\cdot\gamma}{\vdots}}$ such that




for some (necessarily unique) cospan $u \longrightarrow x \longleftarrow v$.


(3) $\Longrightarrow$ (1) Let $\overset{\mathbb{E}}{\underset{\mathbb{B}_{\#}}{p\downarrow}}$ be a discrete fibration with the property of the preceding implication.

We define a small category $\mathbb{X}$ as follows:


— $\mathbb{X}_0 = \{\, E \in \mathbb{E}_0 \mid p(E) = \mathrm{id} \,\}$;
— $\mathbb{X}_1 = \mathbb{E}_0$;
— $\mathrm{dom}, \mathrm{cod} : \mathbb{X}_1 \longrightarrow \mathbb{X}_0$ are the mappings sending $e \in \mathbb{X}_1$ to the unique $\mathrm{dom}(e)$, $\mathrm{cod}(e) \in \mathbb{X}_0$ such that



— $\mathrm{id} : \mathbb{X}_0 \longrightarrow \mathbb{X}_1$ is the inclusion; and
— $_ \cdot _ : \mathbb{X}_1 \underset{\mathbb{X}_0}{\times} \mathbb{X}_1 \longrightarrow \mathbb{X}_1$ is the mapping sending a pair $e, e' \in \mathbb{X}_1$ such that




to the unique $\overset{e\cdot e'}{\underset{p(e)\cdot p(e')}{\vdots}}$ such that




Notice that the above definition only requires pullbacks as in (8) with $g = \mathrm{id}$. The more general pullbacks are needed to establish the associativity law for composition. (We leave the verification of the fact that $\mathbb{X}$ is a category to the reader.)

The mapping $\mathbb{X} \longrightarrow \mathbb{B} : e \longmapsto p(e)$ is functorial (an easy verification, which we also omit) and we denote it f. We claim that f is a ufl functor. Indeed, a morphism $e \in \mathbb{X}_1$ such that $f(e) = \beta_0 \cdot \beta_1$ has a unique factorisation lifting (e_0, e_1) given by the following construction




Finally, we show that $p \cong f_{\#}$ in $\mathbf{DFib}/\mathbb{B}_{\#}$. The isomorphism is given by the functors defined below:

— $\mathbb{E} \longrightarrow \mathbb{X}_{\#} : \epsilon \overset{e}{\downarrow}_{e'} \longmapsto (e_0, e_1) \overset{e}{\downarrow}_{e'}$ where, for $p(e \xrightarrow{\epsilon} e') = (\beta \xrightarrow{(\varphi,\gamma)} \beta')$, we set




— $\mathbb{X}_{\#} \longrightarrow \mathbb{E} : (e_0, e_1) \overset{e}{\downarrow}_{e'} \longmapsto \epsilon \overset{e}{\downarrow}_{e'}$ where we set



□

The property of Lemma 3.7 (3) is an orthogonality condition. Indeed, it states that a presheaf in $\widehat{\mathbb{B}_{\#}}$ is orthogonal to the set of canonical maps $\{ \mathbf{y}(\alpha\cdot\beta) \underset{\mathbf{y}(\beta)}{+} \mathbf{y}(\beta\cdot\gamma) \longrightarrow \mathbf{y}(\alpha\cdot\beta\cdot\gamma) \}$. Moreover, as the diagrams (7) are pullbacks, this property amounts to the sheaf condition

with respect to covers of the form



$$(A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D \text{ in } \mathbb{B}) \tag{10}$$

We denote this family of covers in $\mathbb{B}_{\#}$ by K_3 and write $\mathrm{Ort}(\mathbb{B}_{\#}, \mathrm{K}_3)$ for the full subcategory of $\widehat{\mathbb{B}_{\#}}$ consisting of the presheaves orthogonal (in the above sense) to the covers in K_3. By Proposition 3.6 and Lemma 3.7, in the presence of the axiom of choice, it follows that

$$\mathbf{UFL}/\mathbb{B} \;\simeq\; \mathrm{Ort}(\mathbb{B}_{\#}, \mathrm{K}_3). \tag{11}$$


We thus obtain the following corollary (see Freyd and Kelly (1972), Kelly (1982), Adámek and Rosický (1994) and Borceux (1994)).

Corollary 3.8. $\mathbf{UFL}/\mathbb{B}$ is (equivalent to) a full reflective subcategory of $\widehat{\mathbb{B}_{\#}}$.

One can check that the representables $\mathbf{y}(\beta)$ ($\beta \in |\mathbb{B}_{\#}|$) satisfy the condition of Lemma 3.7 (3). Hence, the Yoneda embedding $\mathbb{B}_{\#} \hookrightarrow \widehat{\mathbb{B}_{\#}}$ factors through the embedding $\mathbf{UFL}/\mathbb{B} \hookrightarrow \widehat{\mathbb{B}_{\#}}$. Explicitly, we obtain the following Yoneda embedding

$$\begin{array}{ccc} \mathbb{B}_{\#} & \xhookrightarrow{\mathbf{u}_{\mathbb{B}}} & \mathbf{UFL}/\mathbb{B} \\ \beta & \longmapsto & \begin{array}{c} {\scriptstyle [\![\beta]\!]} \\ \mathbf{u}(\beta)\downarrow \\ {\scriptstyle \mathbb{B}} \end{array} \end{array} \tag{12}$$


where, for $\beta : B \longrightarrow B'$, the category $[\![\beta]\!]$ has objects given by factorisations $(B \xrightarrow{\varphi} X \xrightarrow{\gamma} B')$ of β in $\mathbb{B}$, morphisms $\chi : (\varphi, X, \gamma) \longrightarrow (\varphi', X', \gamma')$ given by arrows $\chi : X \longrightarrow X'$ in $\mathbb{B}$ such that



(13)

commutes, identities given by $\mathrm{id}_{(\varphi,X,\gamma)} = \mathrm{id}_X$, and composition given as in $\mathbb{B}$; the ufl functor $\mathbf{u}(\beta)$ maps a morphism as in (13) to $X \xrightarrow{\chi} X'$. Following Lawvere (1986c), we will refer to the categories $[\![B \xrightarrow{\beta} B']\!]$ as *interval categories*; they range from the initial object $(\mathrm{id}_B, B, \beta)$ to the terminal object $(\beta, B', \mathrm{id}_{B'})$.

Associated to every functor $f : \mathbb{X} \longrightarrow \mathbb{B}$ there is a natural transformation




where $\overline{f}_x : [\![x]\!] \longrightarrow [\![fx]\!] : (x_0, X, x_1) \longmapsto (fx_0, fX, fx_1)$. This data permits an alternative characterisation of the (UFL) property (*cf.* Lawvere (1986c)):

$$f \text{ is a ufl functor iff } \overline{f} \text{ is a natural isomorphism.}$$

4. Transition graphs

Labelled transition graphs generalise labelled transition systems in that they allow different transitions between states to be labelled by the same action. Specifically, a transition graph consists of a set of states and a set of labelled transitions between states. As an example, we depict below a transition graph over a set of actions A that is not a transition system.



(14)

Morphisms between transition graphs are given by pairs of functions between the underlying sets of states and transitions that respect both the labelling, and the initial and final states of transitions. The category of transition systems is then the full subcategory of the category of transition graphs consisting of objects that are *extensional* in the sense that every transition is uniquely determined by its label, and by its initial and final states.

The category of transition graphs over a set of actions A will be denoted $\mathbf{TG}_A$. The purpose of this section is to analyse *static* and *dynamic* aspects (with emphasis on the latter) of this category. As we will see below, a topos of labelled graphs (see Lawvere (1989)) provides a natural setting for treating static aspects; whilst, as hinted at in Section 2, ufl functors are appropriate structures for the study of the dynamics.

Static aspect. We give the definition of (the static view of) the category of transition graphs.

$$\mathbf{TG}_A \quad = \quad \mathbf{Graph}/\ell(A) \quad \cong \quad \widehat{\mathbf{G}_A}$$

where

$$\mathbf{G}_A{}^{\mathrm{op}} \quad = \quad (\textstyle\int \ell(A))^{\mathrm{op}} \quad \cong \quad \boxed{\mathsf{N} \underset{\mathrm{cod}_a}{\overset{\mathrm{dom}_a}{\leftleftarrows}} \mathsf{E}_a \qquad (a \in A)}$$

Thus, $\mathbf{TG}_A$ is a presheaf topos; the subobject classifier is given by the transition graph (14) equipped with the subobject $\boxed{1 \circlearrowright a \qquad (a \in A)}$ (*cf.* Lawvere (1989) and Bunge and Niefield (1998)). The category of transition systems $\mathbf{TS}_A$ appears as the full subcategory of $\mathbf{TG}_A$ of transition graphs that are *extensional* with respect to

$$\begin{array}{rcll} U \quad = \quad \boxed{\bullet} & \cong & \ell(0 \longrightarrow A) & \text{in } \mathbf{Graph}/\ell(A) \\ & \cong & \mathbf{y}(\mathsf{N}) & \text{in } \widehat{\mathbf{G}_A} \end{array}$$

That is, those transition graphs S with the property that, for all $f, g : G \longrightarrow S$ in $\mathbf{TG}_A$,

$$\text{if } f \circ u = g \circ u, \text{ for all } u : U \longrightarrow G, \text{ then } f = g.$$

It follows that $\mathbf{TS}_A$ is a full reflective exponential ideal of $\mathbf{TG}_A$, and that it is not a topos (in fact, it is the quasitopos of $\neg\neg$-separated objects in $\widehat{\mathbf{G}_A}$). The reflector $\mathbf{TG}_A \longrightarrow \mathbf{TS}_A$ identifies transitions with both the same label, and initial and final states.

Dynamic aspect. We observe that

$$\mathbf{TG}_A \quad \simeq \quad \mathbf{UFL}/\mathbf{M}(A) \tag{15}$$

and argue that this representation embodies the dynamics of transition graphs. The equivalence (15) above for the case $A = 1$ was first noted in Schanuel and Street (1998).

To see (15), observe that the adjunction $\eta, \varepsilon : \mathscr{F} \dashv \mathrm{U} : \mathbf{Cat} \longrightarrow \mathbf{Graph}$ induces the following adjoint situation

$$\mathbf{Graph}/\ell(A) \underset{\Sigma_\eta}{\overset{\eta^*}{\underset{\longrightarrow}{\overset{\longleftarrow}{\top}}}} \mathbf{Graph}/\mathrm{U}\mathscr{F}(\ell A) \underset{\longrightarrow}{\overset{\longleftarrow}{\top}} \mathbf{Cat}/\mathscr{F}(\ell A) \tag{16}$$

where the left adjoint lands in $\mathbf{UFL}/\mathscr{F}(\ell A)$. Moreover, the unit of this adjunction is an isomorphism; whilst, for $f \downarrow_{\mathscr{F}(\ell A)}^{\mathbb{X}}$ in $\mathbf{Cat}/\mathscr{F}(\ell A)$, the component of the counit at f is an isomorphism iff the category $\mathbb{X}$ is freely generated by the morphisms in it over an element of A, iff f is a ufl functor. Thus, the adjunction (16) cuts down to the equivalence (15).

The first dynamic aspect of transition graphs to be considered is the operation that associates a transition graph with its space of *computation paths*. For transition systems, this construction amounts to the well-known coreflection $\mathbf{ST}_A \underset{\longrightarrow}{\overset{\longleftarrow}{\top}} \mathbf{TS}_A^\bullet$ unfolding a transition system with an initial state into a synchronisation tree with root the given initial state (Winskel and Nielsen 1997). The corresponding construction for transition graphs is as follows:

$$\widehat{\mathbf{P}(A)} \underset{\longrightarrow}{\overset{\longleftarrow}{\top}} \mathbf{UFL}/\mathbf{P}(A) \underset{\Sigma_\Delta}{\overset{\Delta^*}{\underset{\longrightarrow}{\overset{\longleftarrow}{\top}}}} \mathbf{UFL}/\mathbf{M}(A) \tag{17}$$

Here the right adjoint unfolds a transition graph into the synchronisation forest of all the possible finite runs of the system. A more conceptual description of this adjunction is *basically* given by the reflection of Corollary 3.8. To see this, for a small category $\mathbb{B}$ let $j : \mathbb{B}_\#^+ \longrightarrow \mathbb{B}_\#$ be the inclusion of the category $\mathbb{B}_\#^+ \cong \sum_{B \in \mathbb{B}_0} B/\mathbb{B}$ of post-extensions in the category $\mathbb{B}_\#$ (see Example 3.5). We have the following adjoint situation

$$\widehat{\mathbb{B}_\#^+} \underset{j_!}{\overset{j^*}{\underset{\longrightarrow}{\overset{\longleftarrow}{\top}}}} \widehat{\mathbb{B}_\#} \underset{\longrightarrow}{\overset{(_)_\#}{\overset{\longleftarrow}{\top}}} \mathbf{UFL}/\mathbb{B}$$

which for $\mathbb{B} = \mathbf{M}(A)$ yields (17).

Finally, we concentrate on process equivalence. A standard such notion is given by *bisimilarity* (Milner 1989). We recall the definition. A *(strong) bisimulation* between transition graphs X and Y is a relation R between their respective state sets such that whenever $x\ R\ y$, then

if $x \xrightarrow[a]{} x'$, then $y \xrightarrow[a]{} y'$ and $x' \; R \; y'$, for some y';

if $y \xrightarrow[a]{} y'$, then $x \xrightarrow[a]{} x'$ and $x' \; R \; y'$, for some x'.

(Here we have used the notation $s \xrightarrow[a]{} s'$ to refer to an anonymous transition from s to s' labelled by a.) One says that two transition graphs with initial state are bisimilar if there is a bisimulation relating the respective initial states.

Following Joyal *et al.* (1996), we provide a treatment of bisimulation for transition graphs via open maps. Very roughly, the idea of this approach is to consider a category of models $\mathscr{M}$ together with a path functor $\pi : \mathscr{P} \longrightarrow \mathscr{M}$, and define the notion of π-bisimilarity as given by a span of π-open maps. A map in $\mathscr{M}$ is said to be π-*open* if the image of every map in $\mathscr{P}$ under π is weakly orthogonal to it; that is, $X \longrightarrow Y$ in $\mathscr{M}$ is π-open if, for every $p : P \longrightarrow Q$ in $\mathscr{P}$, all commutative squares

$$\begin{array}{ccc} \pi P & \longrightarrow & X \\ {\scriptstyle \pi p}\downarrow & \nearrow_{\exists} & \downarrow \\ \pi Q & \longrightarrow & Y \end{array}$$

have a fill-in as indicated.

In the setting of so-called *presheaf models* (Cattani and Winskel 1997), the category of models is given by a presheaf category and the path functor is taken to be the Yoneda embedding. In our context, the structure of the category of paths allows for various alternatives. The straightforward choice is to consider the Yoneda embedding $\mathbf{u}_{\mathbb{B}} : \mathbb{B}_{\#} \hookrightarrow \mathbf{UFL}/\mathbb{B}$ (see (12)); another possibility is its restriction


$$\mathbf{u}^{+}_{\mathbb{B}} : \mathbb{B}^{+}_{\#} \longrightarrow \mathbb{B}_{\#} \hookrightarrow \mathbf{UFL}/\mathbb{B}.$$

For the case under study, $\mathbb{B} = \mathbf{M}(A)$, the path functor $\mathbf{u}^{+}$ is the inclusion $\mathbf{P}(A) \longrightarrow \mathbf{TG}_A$; yielding a situation analogous to the one considered in Joyal *et al.* (1996) for transition systems. It follows that two transition graphs X and Y with initial states $x : U \longrightarrow X$ and $y : U \longrightarrow Y$ are bisimilar iff there exists a span $X \xleftarrow{w_1} W \xrightarrow{w_2} Y$ of $\mathbf{u}^{+}$-open maps such that $(x, y) \in W$ (that is, (x, y) factors through (w_1, w_2)). The notion of bisimulation provided by the path functor $\mathbf{u} : \mathbf{M}(A)_{\#} \longrightarrow \mathbf{TG}_A$ corresponds to *back-and-forth bisimulation* on states (de Nicola *et al.* 1990; Winskel and Nielsen 1997).

5. Path-linearisable categories

In the previous section we regarded and analysed the category of transition graphs $\mathbf{UFL}/\mathbf{M}(A)$ as an *interleaving* model of concurrent computation. In this section control categories like $\mathbf{M}(A)$ are axiomatised as *path-linearisable* categories. Roughly speaking, path-linearisable categories are those in which every commutative square has an essentially unique triangular factorisation (see (CFI) in Definition 5.1 below). Intuitively, a path-linearisable control category is a control category in which any two control paths of morphisms with equal composites can be *linearised* (or *interleaved*) in a canonical fashion,

as exemplified in the following diagram



(18)

where the top and bottom control paths have been linearised (interleaved) as the control path

$$* \longrightarrow \beta_1 \dashrightarrow \alpha_1 \longrightarrow \alpha_2 \dashrightarrow \beta_2 \longrightarrow \beta_3 \dashrightarrow \alpha_3 \longrightarrow \alpha_4 \longrightarrow \bullet.$$

Our main use for path-linearisable categories is in the next section, where we investigate the categories of processes $\mathbf{UFL}/\mathbb{C}$ for a path-linearisable control category $\mathbb{C}$. Thus, from the computational viewpoint, we will be focusing on so-called interleaving models of concurrent processes.


Definition 5.1. A *path-linearisable* category $\mathbb{C}$ is a category satisfying either of the following equivalent conditions.

(CFI) Canonical fill-in.

(C) Cancellation: for every $a \xrightarrow{\alpha} b \underset{\gamma}{\overset{\varphi}{\rightrightarrows}} c \xrightarrow{\beta} d$ in $\mathbb{C}$,

$$\text{if } \alpha \cdot \varphi = \alpha \cdot \gamma \text{ and } \varphi \cdot \beta = \gamma \cdot \beta, \text{ then } \varphi = \gamma\ .$$


(FI) Fill-in: every commutative square



in $\mathbb{C}$ has at least one fill-in as indicated.

(IG) Interval glueing.

For every $a \xrightarrow{\alpha} b \xrightarrow{\beta} c$ in $\mathbb{C}$, the commutative square



(19)

is a pushout in **Cat**.

The equivalence between (CFI) and (IG) will be proved below (Theorem 5.2). Before doing so we mention some examples and comment on the definition.

In a path-linearisable category, fill-in's of commutative squares are canonical in the sense that two parallel fill-in's are equal, whilst two opposite fill-in's are the inverse of


each other. Thus the interval category of a morphism in a path-linearisable category is a linear preorder.

Examples of path-linearisable categories are: free categories on graphs (in particular, the monoid $\mathbf{M}(A)$ for any set A); preorders in which every interval is linear (in particular, the poset $\mathbf{P}(A)$ for any set A); and the additive monoid of non-negative real numbers. These examples are special in that they satisfy the further property:


(II) Every isomorphism is an identity.

This makes the interval categories into linear orders. It follows that the categories satisfying (CFI) and (II) are the *factorisation linearly ordered* categories of Bunge and Niefield (1998).

Examples of path-linearisable categories in which the property (II) does not hold include: groupoids, and the categories



The simplest non-example of a path-linearisable category is the (*independence*) square



which is typical of *non-interleaving* (or *independence*) models of concurrent processes (Winskel and Nielsen 1995).


It was shown in Bunge and Niefield (1998), via model-generated techniques, that, for a category $\mathbb{C}$ satisfying (IG), the category $\mathbf{UFL}/\mathbb{C}$ is a topos. We will give another proof of this result in the next section by exhibiting it as a sheaf topos. We remark that for an arbitrary base category $\mathbb{B}$ the category $\mathbf{UFL}/\mathbb{B}$ need not be a topos. Indeed, this was established by Johnstone for $\mathbb{B}$ the commutative square (Johnstone 1999).

We proceed to show the equivalence between (CFI) and (IG).

Theorem 5.2. The following statements are equivalent.

1 $\mathbb{C}$ has the canonical fill-in property (CFI).

2 For every $a \xrightarrow{\alpha} b \xrightarrow{\beta} c$ in $\mathbb{C}$ the commutative diagram



(20)

is a pushout in **Set**.

3 $\mathbb{C}$ has the interval glueing property (IG).

The implication (3) $\Longrightarrow$ (2) holds because the forgetful functor $(_)_0 : \mathbf{Cat} \longrightarrow \mathbf{Set}$ has a right adjoint and hence preserves colimits.

We prove the implication (2) $\Longrightarrow$ (1). Henceforth we will refer to the statement in Theorem 5.2 (2) as the '$(\mathrm{IG})_0$ property'. We start with the following crucial lemma.

Lemma 5.3. If $\mathbb{C}$ has the $(\mathrm{IG})_0$ property, it satisfies the following property:

Every idempotent is an identity.

Thus it also satisfies the property:

Every retraction is an isomorphism.


Proof. Let e be an idempotent. We write $\approx$ for the equivalence relation on $[\![e]\!]_0 + [\![e]\!]_0$ generated by $\sim$ defined as follows

$$\amalg_1(e \cdot x, y) \sim \amalg_2(x, y \cdot e) \qquad \text{for every } (x, y) \in [\![\mathrm{id}]\!]_0.$$

Notice the important fact that

$$[\amalg_1(e, \mathrm{id})]_{\approx} \quad = \quad \{\, \amalg_1(e, \mathrm{id}), \amalg_2(\mathrm{id}, e) \,\}. \tag{21}$$

We have the following situation



and since $[\![\mathrm{id}, e]\!]_0(e, \mathrm{id}) = (e, e) = [\![\mathrm{id}, e]\!]_0(e, e)$ it follows that $\amalg_1(e, \mathrm{id}) \approx \amalg_1(e, e)$, which, by (21) above, yields $e = \mathrm{id}$. □

It follows that the pushout (20) is simply a union.

Corollary 5.4. Assume that $\mathbb{C}$ has the $(\mathrm{IG})_0$ property. For every $\alpha : a \longrightarrow b$ in $\mathbb{C}$, the functions $[\![\alpha, \mathrm{id}]\!]_0 : [\![\mathrm{id}]\!]_0 \longrightarrow [\![\alpha]\!]_0$ and $[\![\mathrm{id}, \alpha]\!]_0 : [\![\mathrm{id}]\!]_0 \longrightarrow [\![\alpha]\!]_0$ are injective.

Proof. By Lemma 5.3, for every $(x, y) \in [\![\mathrm{id}]\!]_0$, we have that x and y are inverses of each other. Thus, for $(x, y), (x', y') \in [\![\mathrm{id}]\!]_0$, if $[\![\alpha, \mathrm{id}]\!]_0(x, y) = [\![\alpha, \mathrm{id}]\!]_0(x', y')$, then $y = y'$, and so $x = y^{-1} = y'^{-1} = x'$. The injectivity of $[\![\mathrm{id}, \alpha]\!]_0$ is established analogously. □

As a consequence of the above, we have that under the $(\mathrm{IG})_0$ property, for every $a \xrightarrow{\alpha} b \xrightarrow{\beta} c$, the functions $[\![\mathrm{id}, \beta]\!]_0 : [\![\alpha]\!]_0 \longrightarrow [\![\alpha \cdot \beta]\!]_0$ and $[\![\alpha, \mathrm{id}]\!]_0 : [\![\beta]\!]_0 \longrightarrow [\![\alpha \cdot \beta]\!]_0$ are injective. Hence, as we show below, the implication (2) $\Longrightarrow$ (1) holds.

Proposition 5.5. If a category satisfies the $(\mathrm{IG})_0$ property, it also satisfies the canonical fill-in property (CFI).

Proof.

(C) Let $a \xrightarrow{\alpha} b \underset{\gamma}{\overset{\varphi}{\rightrightarrows}} c \xrightarrow{\beta} d$ in $\mathbb{C}$ where $\alpha \cdot \varphi = \alpha \cdot \gamma$ and $\varphi \cdot \beta = \gamma \cdot \beta$. Then we have that φ equals γ because the objects $(\alpha, \varphi), (\alpha, \gamma) \in [\![\alpha \cdot \varphi]\!]_0$ are both mapped under the injection $[\![\mathrm{id}, \beta]\!]_0 : [\![\alpha \cdot \varphi]\!]_0 \longrightarrow [\![\alpha \cdot \varphi \cdot \beta]\!]_0$ to $(\alpha, \varphi \cdot \beta) = (\alpha, \gamma \cdot \beta)$.

(FI) Assume $\alpha \cdot \beta = \varphi \cdot \gamma$. If $(\varphi, \gamma) = [\![\mathrm{id}, \beta]\!]_0(\alpha_0, \alpha_1) \in [\![\alpha \cdot \beta]\!]_0$ for some $(\alpha_0, \alpha_1) \in [\![\alpha]\!]_0$ then α_1 provides the required fill-in; otherwise, $(\varphi, \gamma) = [\![\alpha, \mathrm{id}]\!]_0(\beta_0, \beta_1) \in [\![\alpha \cdot \beta]\!]_0$ for some $(\beta_0, \beta_1) \in [\![\beta]\!]_0$ and the required fill-in is provided by β_0. □

Finally, we show the implication (1) $\Longrightarrow$ (3). The version of Proposition 5.7 below for factorisation linearly ordered categories (categories satisfying (CFI) and (II)) appears in Bunge and Niefield (1998).

Proposition 5.6. If a category satisfies the canonical fill-in property (CFI), it also satisfies the $(\mathrm{IG})_0$ property.

Proof. Let $[\![\alpha]\!]_0 \xrightarrow{f_0} X \xleftarrow{g_0} [\![\beta]\!]_0$ be a cone for the span

$$[\![\alpha]\!]_0 \xleftarrow{[\![\alpha,\mathrm{id}]\!]_0} [\![\mathrm{id}]\!]_0 \xrightarrow{[\![id,\beta]\!]_0} [\![\beta]\!]_0$$

in **Set**.

Since fill-in's are canonical, it follows that the relation $\overset{h_0}{\longmapsto} \subseteq [\![\alpha \cdot \beta]\!]_0 \times X$ given by

$$(\alpha_0, \alpha_1 \cdot \beta) \overset{h_0}{\longmapsto} f_0(\alpha_0, \alpha_1) \quad \text{, for } (\alpha_0, \alpha_1) \text{ in } [\![\alpha]\!]_0$$

$$(\alpha \cdot \beta_0, \beta_1) \overset{h_0}{\longmapsto} g_0(\beta_0, \beta_1) \quad \text{, for } (\beta_0, \beta_1) \text{ in } [\![\beta]\!]_0$$

defines a function $[\![\alpha \cdot \beta]\!]_0 \longrightarrow X$, which can be readily checked to be the required unique mediating morphism. □

Proposition 5.7. If a category satisfies the canonical fill-in property (CFI), it also satisfies the interval glueing property (IG).

Proof. We start with the following lemma: for every $a \xrightarrow{\varphi} b \xrightarrow{\gamma} c$, the functors $[\![\mathrm{id}, \gamma]\!] : [\![\varphi]\!] \longrightarrow [\![\varphi \cdot \gamma]\!]$ and $[\![\varphi, \mathrm{id}]\!] : [\![\gamma]\!] \longrightarrow [\![\varphi \cdot \gamma]\!]$ are embeddings. Indeed, the functor $[\![\mathrm{id}, \gamma]\!]$ is faithful by construction. It is also full because, for objects $(\varphi_0, \varphi_1), (\varphi_0', \varphi_1') \in [\![\varphi]\!]$ and a morphism $\mu : (\varphi_0, \varphi_1 \cdot \gamma) \longrightarrow (\varphi_0', \varphi_1' \cdot \gamma)$ in $[\![\varphi \cdot \gamma]\!]$ we have that $\varphi_0 \cdot \mu = \varphi_0'$, and that $\varphi_1 \cdot \gamma = (\mu \cdot \varphi_1') \cdot \gamma$ and $\varphi_0 \cdot \varphi_1 = \varphi = \varphi_0' \cdot \varphi_1' = \varphi_0 \cdot (\mu \cdot \varphi_1')$. Thus, by cancellation, the equality $\varphi_1 = \mu \cdot \varphi_1'$ holds and we are done. The case for the functor $[\![\varphi, \mathrm{id}]\!]$ is analogous.

It follows that the diagrams (19) consist of injective-on-objects embeddings. We proceed to show that they are pushouts. Let $[\![\alpha]\!] \xrightarrow{f} \mathbb{X} \xleftarrow{g} [\![\beta]\!]$ be a cone for the span

$$[\![\alpha]\!] \xleftarrow{[\![\alpha,\mathrm{id}]\!]} [\![\mathrm{id}]\!] \xrightarrow{[\![\mathrm{id},\beta]\!]} [\![\beta]\!]$$

in **Cat**, and let $h_0 : [\![\alpha \cdot \beta]\!]_0 \longrightarrow \mathbb{X}_0$ be the function given by Proposition 5.6 with respect to the cone $[\![\alpha]\!]_0 \xrightarrow{f_0} \mathbb{X}_0 \xleftarrow{g_0} [\![\beta]\!]_0$ in **Set**.

Notice the following two properties that follow from cancellation.

— For every $(\alpha_0, \alpha_1) \in [\![\alpha]\!]$, $(\beta_0, \beta_1) \in [\![\beta]\!]$ and $\gamma : (\alpha_0, \alpha_1 \cdot \beta) \longrightarrow (\alpha \cdot \beta_0, \beta_1)$ in $[\![\alpha \cdot \beta]\!]$, we have that $\gamma = \alpha_1 \cdot \beta_0$.

— For every $(\alpha_0, \alpha_1) \in [\![\alpha]\!]$, $(\beta_0, \beta_1) \in [\![\beta]\!]$, and $\gamma : (\alpha \cdot \beta_0, \beta_1) \longrightarrow (\alpha_0, \alpha_1 \cdot \beta)$ in $[\![\alpha \cdot \beta]\!]$, we have that β_0 and α_1 are isomorphisms and that $\gamma = {\beta_0}^{-1} \cdot {\alpha_1}^{-1}$.

Thus we may extend the function h_0 to a functor $h = (h_0, h_1) : [\![\alpha \cdot \beta]\!] \longrightarrow \mathbb{X}$ by cases as follows.

1 For $\gamma : (\alpha_0, \alpha_1) \longrightarrow (\alpha_0', \alpha_1')$ in $[\![\alpha]\!]$, set

$$\Big((\alpha_0, \alpha_1 \cdot \beta) \xrightarrow{\gamma} (\alpha_0', \alpha_1' \cdot \beta)\Big) \quad \overset{h_1}{\longmapsto} \quad f_1(\gamma).$$

2 For $\gamma : (\beta_0, \beta_1) \longrightarrow (\beta_0', \beta_1')$ in $[\![\beta]\!]$, set

$$\left((\alpha \cdot \beta_0, \beta_1) \xrightarrow{\gamma} (\alpha \cdot \beta_0', \beta_1')\right) \quad \overset{h_1}{\longmapsto} \quad g_1(\gamma).$$

3 For $(\alpha_0, \alpha_1) \in [\![\alpha]\!]$, $(\beta_0, \beta_1) \in [\![\beta]\!]$, and $\gamma : (\alpha_0, \alpha_1 \cdot \beta) \longrightarrow (\alpha \cdot \beta_0, \beta_1)$ in $[\![\alpha \cdot \beta]\!]$, set

$$\gamma \quad \overset{h_1}{\longmapsto} \quad f_1\left((\alpha_0, \alpha_1) \xrightarrow{\alpha_1} (\alpha, \mathrm{id})\right) \cdot g_1\left((\mathrm{id}, \beta) \xrightarrow{\beta_0} (\beta_0, \beta_1)\right).$$


4 For $(\alpha_0, \alpha_1) \in [\![\alpha]\!]$, $(\beta_0, \beta_1) \in [\![\beta]\!]$, and $\gamma : (\alpha \cdot \beta_0, \beta_1) \longrightarrow (\alpha_0, \alpha_1 \cdot \beta)$ in $[\![\alpha \cdot \beta]\!]$, set

$$\gamma \quad \overset{h_1}{\longmapsto} \quad g_1\left((\beta_0, \beta_1) \xrightarrow{{\beta_0}^{-1}} (\mathrm{id}, \beta)\right) \cdot f_1\left((\alpha, \mathrm{id}) \xrightarrow{{\alpha_1}^{-1}} (\alpha_0, \alpha_1)\right).$$

It is easy to check that h is well-defined and has the required universal property. □

We conclude this section with a property of twisted arrow categories of categories with the cancellation property that gives insight into these notions. Intuitively, the property says that a path appearing in two non-overlapping parts of another path is a deformation.

Proposition 5.8. For $\mathbb{C}$ satisfying the cancellation property, every commutative diagram



in $\mathbb{C}_{\#}$ is such that ρ and σ_1 are isomorphisms. Hence, the map $(\sigma_0, \sigma_1) : \rho \longrightarrow \alpha \cdot \beta$ factors through $(\alpha, \mathrm{id}) : \beta \longrightarrow \alpha \cdot \beta$.


Proof. We have that $\varphi \cdot \alpha \cdot \beta = \varphi \cdot \alpha \cdot \beta \cdot (\gamma_0 \cdot \tau_0 \cdot \rho \cdot \sigma_1)$ and that $(\gamma_0 \cdot \tau_0 \cdot \rho \cdot \sigma_1) \cdot \gamma_0 \cdot \gamma \cdot \gamma_1 = \gamma_0 \cdot \gamma \cdot \gamma_1$, and so, by cancellation, $\gamma_0 \cdot \tau_0 \cdot \rho \cdot \sigma_1 = \mathrm{id}$. Since in a category satisfying the cancellation property retractions are isomorphisms, it follows that γ_0, τ_0, ρ, σ_1 are isos. The map (σ_0, σ_1) factors as $(\beta \cdot {\sigma_1}^{-1} \cdot \rho^{-1}, \sigma_1) \cdot (\alpha, \mathrm{id})$. □

6. Sheaf models of linearly-controlled processes

We study the categories $\mathbf{UFL}/\mathbb{C}$ for $\mathbb{C}$ a path-linearisable category. The fact that the control category is path-linearisable leads to considering processes with a category of states in which any two evolution paths with equal endpoints can be linearised into one. Indeed, we have the following proposition.

Proposition 6.1. For a ufl functor $f \downarrow \begin{matrix}\mathbb{X}\\ \mathbb{C}\end{matrix}$ over a path-linearisable category $\mathbb{C}$, the category $\mathbb{X}$ is path-linearisable.


Proof. The proof follows from the commutativity of the following cube



for every $X \xrightarrow{x} Y \xrightarrow{y} Z$ in $\mathbb{X}$. □

As a consequence, interval categories in a path-linearisable category are path linearisable (see (12)).

We proceed to show that the categories $\mathbf{UFL}/\mathbb{C}$ for $\mathbb{C}$ a path-linearisable category are sheaf toposes – they may be regarded as sheaf models of linearly-controlled processes. Our approach is to establish the equivalence between the orthogonality condition of Lemma 3.7 (3) and a sheaf condition. To this end, we need to consider binary versions of the covers K_3 introduced in (10). For a small category $\mathbb{C}$, let K_2 be the family of covers



$(A \xrightarrow{\alpha} Y \xrightarrow{\gamma} D \text{ in } \mathbb{C})$

in $\mathbb{C}_{\#}$. These covers need not be pretopologies; however, the stability axiom can be easily guaranteed.


Proposition 6.2. For a small category $\mathbb{C}$, the family of covers K_2 satisfies the stability axiom for a pretopology iff the category $\mathbb{C}$ has the fill-in property (FI).

Proof.


($\Longrightarrow$) Let $\alpha \cdot \beta = \varphi \cdot \gamma$. Consider the cover $\alpha \xrightarrow{(\mathrm{id},\beta)} \varphi \cdot \gamma \xleftarrow{(\alpha,\mathrm{id})} \beta$ and the map $(\varphi, \gamma) : \mathrm{id} \longrightarrow \varphi \cdot \gamma$. There exists a cover $\mu \xrightarrow{(\mathrm{id},\nu)} \mathrm{id} \xleftarrow{(\mu,\mathrm{id})} \nu$ such that, for instance, the composite $\mu \xrightarrow{(\mathrm{id},\nu)} \mathrm{id} \xrightarrow{(\varphi,\gamma)} \varphi \cdot \gamma$ factors through $\alpha \xrightarrow{(\mathrm{id},\beta)} \varphi \cdot \gamma$ or through $\beta \xrightarrow{(\alpha,\mathrm{id})} \varphi \cdot \gamma$. In the first case, if $(\mathrm{id}, \nu) \cdot (\varphi, \gamma) = (\rho, \sigma) \cdot (\mathrm{id}, \beta)$, then $\mu \cdot \sigma$ is a fill-in for the equal pair of arrows $\alpha \cdot \beta$ and $\varphi \cdot \gamma$. In the second case, if $(\mathrm{id}, \nu) \cdot (\varphi, \gamma) = (\rho, \sigma) \cdot (\alpha, \mathrm{id})$, then ρ provides the fill-in.

($\Longleftarrow$) Consider the cover $\alpha \xrightarrow{(\mathrm{id},\beta)} \alpha \cdot \beta \xleftarrow{(\alpha,\mathrm{id})} \beta$ and let $(\mu, \nu) : \gamma \longrightarrow \alpha \cdot \beta$. The proof proceeds by inspecting the linearisations (interleavings) of the equal pair of arrows $\alpha \cdot \beta$

and $\mu \cdot \gamma \cdot \nu$. The only non-trivial situation is



in which case we have the factorisations below



□

Let $\overline{\mathrm{K}_2}$ be the closure of K_2 under transitivity. Explicitly, the covers of $\overline{\mathrm{K}_2}$ are given by the families

$$\{(\gamma_1 \cdot \ldots \cdot \gamma_{i-1}, \gamma_{i+1} \cdot \ldots \cdot \gamma_n) : \gamma_i \longrightarrow \gamma_1 \cdot \ldots \cdot \gamma_i \cdot \ldots \cdot \gamma_n \mid 1 \leqslant i \leqslant n\}.$$

As K_2 satisfies the identity axiom of pretopologies, for a category $\mathbb{C}$ with the fill-in property, we obtain a site $(\mathbb{C}_{\#}, \overline{\mathrm{K}_2})$. Now, following Johnstone (see Johnstone (1999)), we restrict attention to categories satisfying the cancellation property. We thus consider path-linearisable categories. For them we show that the sheaf condition with respect to $\overline{\mathrm{K}_2}$ is equivalent to orthogonality with respect to K_3. This, by (11) above, establishes Theorem 6.4.

Proposition 6.3. Let $\mathbb{C}$ be a path-linearisable category. For a presheaf in $\widehat{\mathbb{C}_{\#}}$ the following statements are equivalent.

1 The presheaf is orthogonal to every cover in K_3.
2 The presheaf is orthogonal to every cover in K_2.
3 The presheaf is a $\overline{\mathrm{K}_2}$-sheaf.

Proof.
(1) $\Longrightarrow$ (2) This holds vacuously.

(2) $\Longrightarrow$ (3) For a presheaf P in $\widehat{\mathbb{C}_{\#}}$ we show by induction on the size of the covers in $\overline{\mathrm{K}_2}$ that P satisfies the sheaf condition.

For $n = 1$ the sheaf condition is trivial, and for $n = 2$ it amounts to orthogonality with respect to covers in K_2. For the inductive step, assume that P satisfies the sheaf condition for all covers in $\overline{\mathrm{K}_2}$ of size n ($n \geqslant 2$). Consider a cover

$$\{\gamma_i \longrightarrow \gamma_1 \cdot \ldots \cdot \gamma_i \cdot \ldots \cdot \gamma_{n+1} \mid 1 \leqslant i \leqslant n+1\} \tag{22}$$

and let

$$\{x_i : \gamma_i \longrightarrow P \mid 1 \leqslant i \leqslant n+1\} \tag{23}$$

be a matching family for it. Define $x_{1,2} : \gamma_1 \cdot \gamma_2 \longrightarrow P$ as the unique amalgamation, given


by orthogonality with respect to K_2, of the matching family $\gamma_1 \xrightarrow{x_1} P \xleftarrow{x_2} \gamma_2$ for the cover $\gamma_1 \longrightarrow \gamma_1 \cdot \gamma_2 \longleftarrow \gamma_2$ in K_2. Observe that the family

$$\{ x_{1,2} \} \cup \{ x_i \mid 3 \leqslant i \leqslant n+1 \} \tag{24}$$

is a matching family for the cover

$$\{ \gamma_1 \cdot \gamma_2 \longrightarrow \gamma_1 \cdot \gamma_2 \ldots \cdot \gamma_{n+1} \} \cup \{ \gamma_i \longrightarrow \gamma_1 \cdot \ldots \cdot \gamma_i \cdot \ldots \cdot \gamma_{n+1} \mid 3 \leqslant i \leqslant n+1 \} \tag{25}$$

in $\overline{\mathrm{K}_2}$ because, by Proposition 5.8, whenever we have a common restriction as below



$(3 \leqslant i \leqslant n+1)$

the map $\rho \longrightarrow \gamma_1 \cdot \gamma_2$ factors through $(\gamma_1, \mathrm{id}) : \gamma_2 \longrightarrow \gamma_1 \cdot \gamma_2$. By the induction hypothesis, there exists a unique amalgamation $x : \gamma_1 \cdot \ldots \cdot \gamma_{n+1} \longrightarrow P$ such that $(\gamma_1 \cdot \gamma_2 \longrightarrow \gamma_1 \cdot \gamma_2 \cdot \gamma_3 \cdot \ldots \cdot \gamma_{n+1}) \cdot x = x_{1,2}$ and $(\gamma_i \longrightarrow \gamma_1 \cdot \ldots \cdot \gamma_{n+1}) \cdot x = x_i$ $(3 \leqslant i \leqslant n+1)$. Clearly, x is an amalgamation for the matching family (23) with respect to the cover (22). It is unique because any such amalgamation is also an amalgamation for the matching family (24) with respect to the cover (25).

(3) $\Longrightarrow$ (1) Let P be a $\overline{\mathrm{K}_2}$-sheaf.

For $A \xrightarrow{\alpha} B \xrightarrow{\beta} C \xrightarrow{\gamma} D$ in $\mathbb{C}$, let $\alpha \cdot \beta \xrightarrow{x} P \xleftarrow{y} \beta \cdot \gamma$ be a matching family for the cover $\alpha \cdot \beta \longrightarrow \alpha \cdot \beta \cdot \gamma \longleftarrow \beta \cdot \gamma$ in K_3, and let $\{ u : \alpha \longrightarrow P,\ v : \beta \longrightarrow P,\ w : \gamma \longrightarrow P \}$ be a matching family for the cover $\{ \alpha \longrightarrow \alpha \cdot \beta \cdot \gamma,\ \beta \longrightarrow \alpha \cdot \beta \cdot \gamma,\ \gamma \longrightarrow \alpha \cdot \beta \cdot \gamma \}$ in $\overline{\mathrm{K}_2}$. The result follows form the two observations below:

1 The identities $u = (\mathrm{id}, \beta) \cdot x$, $v = (\alpha, \mathrm{id}) \cdot x = (\mathrm{id}, \gamma) \cdot y$, and $w = (\beta, \mathrm{id}) \cdot y$ hold if and only if x is the unique amalgamation of the matching family $\{ u, v \}$ with respect to the cover $\alpha \longrightarrow \alpha \cdot \beta \longleftarrow \beta$ in $\overline{\mathrm{K}_2}$ and y is the unique amalgamation of the matching family $\{ v, w \}$ with respect to the cover $\beta \longrightarrow \beta \cdot \gamma \longleftarrow \gamma$ in $\overline{\mathrm{K}_2}$.

2 The amalgamations of the matching family $\alpha \cdot \beta \xrightarrow{x} P \xleftarrow{y} \beta \cdot \gamma$ with respect to the cover $\alpha \cdot \beta \longrightarrow \alpha \cdot \beta \cdot \gamma \longleftarrow \beta \cdot \gamma$ in K_3 are in bijective correspondence with the amalgamations of the matching family $\{ (\mathrm{id}, \beta) \cdot x,\ (\alpha, \mathrm{id}) \cdot x = (\mathrm{id}, \gamma) \cdot y,\ (\beta, \mathrm{id}) \cdot y \}$ with respect to the cover $\{ \alpha \longrightarrow \alpha \cdot \beta \cdot \gamma,\ \beta \longrightarrow \alpha \cdot \beta \cdot \gamma,\ \gamma \longrightarrow \alpha \cdot \beta \cdot \gamma \}$ in $\overline{\mathrm{K}_2}$. □

Theorem 6.4. For $\mathbb{C}$ a path-linearisable category, we have the following equivalence:

$$\mathbf{UFL}/\mathbb{C} \quad \simeq \quad \underline{\mathrm{Sh}}(\mathbb{C}_{\#}, \overline{\mathrm{K}_2}).$$

7. Concluding remarks

Our treatment of categories of processes, we believe, provides the basis for a fruitful approach to a conceptual analysis of general processes with state. From this viewpoint this paper opens up new possibilities, and we consider it as ground work for future

developments and refinements of some of the topics dealt with here. In particular, investigations related to hybrid systems will be pursued elsewhere.

The fact that $\mathbf{UFL}/\mathbb{C}$ is a topos (for $\mathbb{C}$ path linearisable) has not been exploited here. Nor has its feature of 'single universe' (for discrete fibrations and discrete opfibrations) been analysed. This relates to considering processes with 'past' and 'future', or with history.

The general considerations of this paper make sense not just for categories and functors, but also in other contexts; for example, for multicategories and multifunctors (Lambek 1989) and for computads (Street 1996).

References

Adámek, J. and Rosický, J. (1994) *Locally Presentable and Accessible Categories*, London Mathematical Society Lecture Note Series **189**, Cambridge University Press.

Bunge, M. and Niefield, S. (1998) Exponentiability and single universes. (To appear in *J. Pure Appl. Algebra*.)

Borceux, F. (1994) *Handbook of Categorical Algebra*, Number 50–52 in Encyclopedia of Mathematics and its Applications, Cambridge University Press.

Cattani, G. L. and Winskel, G. (1997) Presheaf models for concurrency. In: van Dalen, D. and Bezem, M. (eds.) Proceedings of the CSL'96 Conf. *Springer-Verlag Lecture Notes in Computer Science* **1258** 58–75.

de Nicola, R., Montanari, U. and Vaandrager, F. (1990) Back and forth bisimulations. In: Proceedings of the CONCUR'90 Conf. *Springer-Verlag Lecture Notes in Computer Science* **458** 152–165.

Freyd, P. J. and Kelly, G. M. (1972) Categories of continuous functors I. *J. of Pure Appl. Algebra* **2** 169–191. (Erratum: *J. Pure Appl. Algebra* (1974) **4** 121).

Joyal, A., Nielsen, M. and Winskel, G. (1996) Bisimulation from open maps. *Inform. and Comput.* **127** (2) 164–185.

Johnstone, P. T. (1999) A note on discrete Conduché fibrations. *Theory and Applications of Categories* **5** 1–11.

Kelly, G. M. (1982) *Basic concepts of enriched category theory*, London Mathematical Society Lecture Note Series **64**, Cambridge University Press.

Lambek, J. (1989) Multicategories revisited. In: Categories in Computer Science and Logic. *Contemporary Mathematics* **92**, AMS 217–239.

Lawvere, F. W. (1986a) Categories of spaces may not be generalized spaces as exemplified by directed graphs. *Rev. Colombiana Mat.* **20** 179–185.

Lawvere, F. W. (1986b) Introduction. In: Lawvere, F. W. and Schanuel, S. H. (eds.) Categories in continuum physics. *Springer-Verlag Lecture Notes in Computer Science* **1174** 1–16.

Lawvere, F. W. (1986c) State categories and response functors. (Unpublished, May 1986.)

Lawvere, F. W. (1989) Qualitative distinctions between some toposes of generalized graphs. In: Categories in Computer Science and Logic. *Contemporary Mathematics* **92**, AMS 261–299.

Milner, R. (1989) *Communication and Concurrency*, Prentice Hall.

Mac Lane, S. (1971) *Categories for the Working Mathematician*, Springer-Verlag.

Mac Lane, S. and Moerdijk, I. (1994) *Sheaves in geometry and logic: A first introduction to topos theory*, Universitext, Springer-Verlag.

Schanuel, S. and Street, R. (1988) The length of a pasting diagram. (Unpublished, June 1988.)

Street, R. (1996) Categorical structures. In: *Handbook of algebra* **1**, North-Holland 529–577.

Winskel, G. and Nielsen, M. (1995) Models for concurrency. In: *Handbook of logic in computer science* **4**, Oxford Univ. Press 1–148.

Winskel, G. and Nielsen, M. (1997) Presheaves as transition systems. In: Partial order methods in verification. *DIMACS Ser. Discrete Math. Theoret. Comput. Sci.* **29** 129–140.